

# Antimicrobial activity of selected red and white wines against *Escherichia coli*: In vitro inhibition using fish as food matrix

---

**Wheeler [Santoro], Heidi Christine**

**Master's thesis / Diplomski rad**

**2020**

*Degree Grantor / Ustanova koja je dodijelila akademski / stručni stupanj:* **University of Split, School of Medicine / Sveučilište u Splitu, Medicinski fakultet**

*Permanent link / Trajna poveznica:* <https://um.nsk.hr/um:nbn:hr:171:870221>

*Rights / Prava:* [In copyright](#)/[Zaštićeno autorskim pravom.](#)

*Download date / Datum preuzimanja:* **2024-12-19**



*Repository / Repozitorij:*

[MEFST Repository](#)



**UNIVERSITY OF SPLIT  
SCHOOL OF MEDICINE**

**HEIDI CHRISTINE SANTORO**

**ANTIMICROBIAL ACTIVITY OF SELECTED RED AND  
WHITE WINES AGAINST ESCHERICHIA COLI: IN VITRO  
INHIBITION USING FISH AS FOOD MATRIX**

**Diploma thesis**

**Academic year: 2019/2020**

**Mentor: Prof. Mladen Boban, MD, PhD**

**Split, July 2020**

## TABLE OF CONTENTS

1. INTRODUCTION & RESEARCH OBJECTIVES .....	1
2. MATERIALS & METHODS.....	5
2.1. Wine Samples and Technology .....	6
2.2. Wine Chemistry Analyses .....	6
2.3. Analysis of Total Phenolics, Tannins and Anthocyanins .....	7
2.4. The Escherichia coli Strains and Inoculum Preparation .....	7
2.5. In Vitro Evaluation of Antimicrobial Activity of Wines.....	8
2.5.1. Agar-Disk and Agar-Well Diffusion Assays .....	8
2.5.2. Minimal Inhibitory Concentration (MIC) .....	8
2.6. Preparation of Fish Samples and Inoculation with E. coli .....	9
2.6.1. Enumeration of <i>E. coli</i> .....	9
2.6.2. pH Analyses .....	10
2.7. Statistical Analyses.....	10
3. RESULTS & DISCUSSION .....	11
3.1. Wine Chemistry .....	12
3.2. Antimicrobial Activity.....	12
3.3. The Experiment on Fish Samples .....	15
4. CONCLUSIONS .....	19
5. REFERENCES .....	21
6. SUMMARY .....	26
7. SAŽETAK.....	28
8. CURRICULUM VITAE .....	30

## ACKNOWLEDGEMENTS

*My gratitude and thanks go out to those who have been a part of my journey the past six years. In particular I wish to say a very big thank you to my mentor, Prof. dr. sc. Mladen Boban for introducing me to the medial world of wine and helping me along the way, as well as Assoc. Prof Vida Šimat and Assist. Prof Danijela Skroza, both of whom I could not have done this without. I thank my son, Oliver, for joyfully going through everything alongside me.*

*This thesis has been published as an original scientific article in MDPI Foods journal (Q1, IF: 4.092): Santoro, H-C.; Skroza, D., Dugandžić, A.; Boban, M.; Šimat, V. (2020). Antimicrobial activity of selected red and white wines against Escherichia coli: in vitro inhibition using fish as food matrix. Foods, 9, 936; doi:10.3390/foods9070936.*

*This research was funded by the Croatian Science Foundation (Project No. 8652).*

## **1. INTRODUCTION & RESEARCH OBJECTIVES**

Food-borne bacteria have become an ever-increasing challenge, in both commercial and domestic use. Contaminated food continues to cause millions of people to fall ill, with many succumbing to fatality due to foodborne bacterial infections [1]. *Escherichia coli* are among the most common food-borne pathogens related to food spoilage and food poisoning. *E. coli* are found in the environment and the lower intestines of warm-blooded organisms as a part of normal microflora and represent mostly harmless bacteria. Certain types of *E. coli*, such as shiga-like-toxin-producing *E. coli* (serotype O157:H7, SLTEC), contaminate different types of foods and water, presenting a risk for humans [2,3]. The main sources of human infections are undercooked and contaminated meat and fish, as well as dairy products made from raw milk. Despite various measures taken during processing, consumers can still be exposed to this pathogen by consumption and handling of improperly prepared foods or through cross-contamination as another major risk factor [3,4].

*E. coli* can tolerate and rapidly adapt to a variety of stressful conditions such as fluctuations and extremes of temperature, high osmolarity, and low pH [5]. One potential approach to reducing the counts of *E. coli* could be the application of natural and sustainable substances with antimicrobial properties. Red and white wines have demonstrated antimicrobial activity under various experimental conditions against spoilage and pathogenic food bacteria such as *Bacillus* spp., *Escherichia* spp., *Listeria* spp., *Salmonella* spp., *Sigella*, *Pseudomonas* spp., *Campylobacter jejuni*, *Yersinia enterocolitica*, *Vibrio* spp., and *Staphylococcus aureus* [6–19].

It is generally held that wine's biological potential is related to a large number of polyphenolic compounds [12]. Polyphenols (phenolics) are plant secondary metabolites, which have essential roles in plant physiology but have potential health benefits on the human organism, mainly as antioxidant, anti-allergic, anti-inflammatory, anticancer, antihypertensive, and antimicrobial agents [20]. The antimicrobial properties of polyphenols have been widely studied to propose new methods of food preservation and enhance food safety without the use of synthetic additives. Among polyphenol classes, *E. coli* showed sensitivity to phenolic acids, hydrolysable and condensed tannins, flavan-3-ols, and flavonols. All of these are found in fruits and vegetables, and plant-derived beverages, such as tea and wine [20]. Red and white wines contain a variety of phenolic compounds that have been

associated with numerous health benefits mentioned earlier but have also been active against foodborne pathogens.

In past years, great attention has been paid to wine polyphenols and naturally produced macerated white wines are becoming increasingly popular; however their biological effects are rarely studied. These wines have a high content of biologically active compounds and among them; polyphenols are considered the main factor responsible for particular biological properties. The phenolic composition of wines varies with a grape variety, but it is also influenced by viticultural and environmental factors, and the wine-making process. However, the skins and seeds of grapes are known to be rich sources of polyphenolic compounds, both flavonoids and non-flavonoids [21], and thus phenolic-rich wines are commonly obtained through maceration step, in which grape juice is left in contact with grape skins, seeds and even stems, for weeks and even months [22]. Although wine polyphenols have been intensively studied, it has been demonstrated that the synergistic effect of phenolics with other wine components, such as low pH, ethanol, and organic acids, remains the key mechanism of wine bioactivity [11,12,23]. The specific components of wine and their relative contribution to its antimicrobial activity are still a matter of debate.

There have been a limited number of studies investigating the antimicrobial effect of wine used in the process of food preparation, either as an ingredient or marinade. In available studies, the wine was not tested intact, but in the form of wine extract or in the combination with soya sauce, oregano and thyme essential oils, garlic juice, or even pure phenolic compounds [7–10,24,25]. Marinade recipes containing red or white wine enriched with plant essential oils demonstrated strong activity under laboratory conditions against major foodborne pathogens [11]. Also, the wine application in foods contributed to the prolongation of shelf life of meat [8] and fish [24] products with the ability to reduce total viable count in the products. Both red and white wines from different regions and grape varieties showed the antimicrobial potential against *E. coli*, however, the in vitro effect was not often confirmed in food models [12,19].

There is an increasing interest in investigations into the use of bioactive natural compounds for the enhancement of food safety and product quality. Microbiological challenge tests are an important tool necessary for simulation of assessment of real-life food safety and prediction of safety risks. The use of natural antimicrobials and preservative sources in foods such as wine would help reduce the use of synthetic additives and aggressive processing, without compromising food safety. In this regard, the main objective of the present study was


to test the antimicrobial activity of five wines against two *E. coli* strains and to verify the findings in wine-marinated fish fillets as a food matrix.



## **2. MATERIALS & METHODS**

## 2.1. Wine Samples and Technology

The five wines, three white wines produced from the grape variety Graševina in 2015 at the Krauthaker winery, and one red and one rosé produced from the grape variety Plavac Mali in 2015 at the Volarević winery, Croatia, were tested in vitro for antimicrobial activity in the standard growing media and in fish fillets as food matrix. The technology of production of the wines tested is schematically shown in Figure 1. The standard white wine (GS) and rosé wine (PM 0) were produced following a classic methodology requiring grape juice to be separated from the hard parts of the grapes during fermentation, without maceration. Conversely, the Georgian wine production principles were used in obtaining macerated white wines rich in polyphenols. Following spontaneous fermentation, and without removing grape seeds and skins, the tanks were sealed air-tight, allowing the wine further elaboration for four months. Part of the wine so produced was bottled without further treatment (GM), while sulphur (GM + SO<sub>2</sub>) was added to the other part as a preservative. The red wine (PM 7) production process included both fermentation and maceration for seven days in contact with the grape seeds and skins (Figure 1).



**Figure 1.** Technology of the tested wines.

## 2.2. Wine Chemistry Analyses

The enological analysis of all wines included ethanol and sulfur content, along with pH and acidity values. These were determined according to the Compendium of International Methods of Analysis of Wines and Musts of the International Organisation of Vine and Wine (OIV), and are shown in Table 1.

**Table 1.** Chemical composition of tested wines.

Wine Samples *	pH	Acidity (g/L) **	Alcohol (%)	Total Phenolics (mg GAE/L)	Anthocyanins (mg/L)	Tannins (mg/L)	Free SO <sub>2</sub> (mg/L)	Total SO <sub>2</sub> (mg/L)
GS	3.54	4.9	13.0	305 <sup>±</sup> 3.5 <sup>a</sup>	0 a	0.26 <sup>±</sup> 0.01 <sup>a</sup>	27	124
GM	3.94	4.8	13.3	2699 <sup>±</sup> 8.2 <sup>b</sup>	0 b	3.34 <sup>±</sup> 0.10 <sup>b</sup>	1	3
GM + SO <sub>2</sub>	3.90	5.4	12.8	2850 <sup>±</sup> 34.6 <sup>b</sup>	0 c	4.93 <sup>±</sup> 0.19 <sup>c</sup>	5	99
PM 0	3.54	3.8	14.3	539 <sup>±</sup> 22.2 <sup>c</sup>	1.97 <sup>±</sup> 0.04 <sup>b</sup>	1.08 <sup>±</sup> 0.07 <sup>d</sup>	9	173
PM 7	3.61	5.7	13.7	1988.96 <sup>±</sup> 35.0 <sup>d</sup>	30.14 <sup>±</sup> 6.52 <sup>d</sup>	4.93 <sup>±</sup> 0.46 <sup>e</sup>	6	89

\* GS—standard white wine, GM and GM + SO<sub>2</sub>—macerated white wine with and without SO<sub>2</sub>, PM 0—rosé, PM 7— macerated red wine: \*\* expressed as tartaric acid; <sup>a-e</sup> mean value ± standard deviation in the same column followed by different superscript are significantly different ( $p < 0.05$ ).

### 2.3. Analysis of Total Phenolics, Tannins and Anthocyanins

The total phenolic content of the wine samples was measured spectrophotometrically using the Folin-Ciocalteu method [21,26] with the results expressed as mg of gallic acid equivalents per liter (mg GAE/L).

The total tannin contents were measured by acidic hydrolysis of proanthocyanidins resulting in carbocation formation with partial conversion into red cyanidin using the method of Ribéreau-Gayon and Stonestreet [27] and concentration was obtained in g/L.

Anthocyanin content was determined by the SO<sub>2</sub> bleaching procedure and results are expressed as mg/L [28].

All measurements were carried out in triplicate using a SPECORD 200 Plus (Analytik Jena AG, Jena, Germany).

### 2.4. The *Escherichia coli* Strains and Inoculum Preparation

Bacterial strains *E. coli* WDCM00012 (ATCC8739, *E. coli* 12) and WDCM00013 (ATCC25922, *E. coli* 13) were used as test microorganisms. Determination of antimicrobial activity was performed using Müller Hinton Broth/Agar (MHB, MHA; Biolife, Italy). The cell culture suspension was prepared by selection of a colony from the 24-hr old MHA plates, with

adjustment in MHB to include a concentration of approximately  $10^8$  colony forming units per mL (CFU/mL) according to the turbidity of 0.5 McFarland scale standard (Densomat, bioMerieux, Marcy l'Etoile, France). An aliquot of 2 mL of cell culture suspension was diluted with 38 mL of MHB medium for inoculum preparation (initial concentration  $10^7$  CFU/mL).

## **2.5. In Vitro Evaluation of Antimicrobial Activity of Wines**

### **2.5.1. Agar-Disk and Agar-Well Diffusion Assays**

The protocols used for agar-disk and agar-well diffusion assays are defined by Clinical and Laboratory Standards Institute (CLSI) [29,30]. The Petri plates containing 20 mL MHA were inoculated by spreading microbial inoculum over the entire agar surface using sterile cotton swabs rolled in the inoculum. For the disk diffusion method, filter paper discs of 6 mm in diameter containing 25  $\mu$ L of the wine samples were placed on the agar surface. The same was done for the antibiotics, chloramphenicol, gentamicin, and tigecycline, used as control. The Petri plates were incubated at 37 °C for 20–24 h. The inhibition zone of the antibiotics was in the range proposed by CLSI and the European Committee on Antimicrobial Susceptibility Testing (EUCAST) [30,31] for both *E. coli* strains.

For agar-well diffusion assays, four equidistant holes were made in the agar using sterile cork borers ( $\varnothing = 7$  mm) and 50  $\mu$ L of the wine samples was added directly in the wells. The agar plates were at 4 °C for one hour, following incubation at 37 °C for 20–24 h. The inhibition zones formed on the medium were measured in millimeters (mm) [32]. If the inhibition zone was  $\geq 12$  mm, the tested sample was considered to have good inhibitory effect [33]. All measurements were done in triplicate with two positive growth controls included (ethanol in the concentration of 13–15%).

### **2.5.2. Minimal Inhibitory Concentration (MIC)**

Minimal inhibitory concentration was measured according to the previously described method by [32,34]. Shortly 50  $\mu$ L of bacterial inoculum and 50  $\mu$ L of two-fold serially diluted wine samples were added to the wells of a sterile 96-well microtitre plate. Three antibiotics (chloramphenicol, gentamicin, and tigecycline) and ethanol (13–15% vol) were used as control. The MIC values of chloramphenicol, gentamicin, and tigecycline were in the range proposed by CLSI and EUCAST standards, 4, 0.5 and 0.15  $\mu$ g/mL, respectively [30,31]. The MIC was defined as the lowest concentration displaying the ability to inhibit any visible bacterial growth

after adding 10  $\mu\text{L}$ /well of INT (2-p-iodophenyl-3-p-nitrophenyl-5-phenyl tetrazolium chloride, Sigma, Darmstadt, Germany).

All measurements of MIC values were done in triplicate and expressed as mg GAE/L.

## **2.6. Preparation of Fish Samples and Inoculation with *E. coli***

Seabass (*Dicentrarchus labrax*) were purchased at the local fish market and transported to the lab packed with ice in polystyrene boxes. The fish were of class L, weighing ~400–600 g each. All further preparations were performed under sterile conditions. The fish was filleted and the fillets thoroughly washed in sterile distilled water. Five groups of fish fillets, each weighing approximately 200 g, were marinated in a mixture of wine and sterile distilled water (400 mL) prepared in the ratio 1:1. The control sample was kept in sterile distilled water. After two hours, each group of fillets was drained and ground in a knife mill down to 40–50 mm particles (GRINDOMIX GM 200, Retsch GmbH, Haan, Germany). Next, 50 g of ground meat was separated for pH measurement.

For the challenge test experiment, triplicate packages containing chunks of  $25 \pm 0.01$  g of ground wine-marinated fish meat were placed in sterile vacuum bags, inoculated with *E. coli* strains (0.5 mL of liquid inoculum prepared as described in Section 2.4) to achieve an initial value of  $\sim 7$  log CFU/ 25 g and vacuum sealed. The samples were kept at room temperature for 10 min to allow possible attachment and diffusion [7] and then stored at  $4 \pm 1$  °C. The number of *E. coli* was enumerated after three and seven days. In addition, triplicate packages of wine-marinated fish meat without inoculum were tested for *E. coli*.

### **2.6.1. Enumeration of *E. coli***

The enumeration of *E. coli* was done using ISO methodology [35]. Briefly, the samples were transferred to stomacher bags with 225 mL of sterile peptone water and homogenized using a laboratory blender (Stomacher, Maxicator, IUL S.A., Barcelona, Spain) for three minutes. The serial of decimal dilutions was prepared and distributed to sterile Petri dish followed by the Tryptone Bile X-GLUC agar (Biolife, Milan, Italy) and incubated at 44 °C for 24–48 h. Plates with 10 to 150 blue-green colonies were counted. Confirmation of *E. coli* was done by Colilert-18 (Quanti-Tray/2000) method. According to the manufacturer's instructions, yellow wells indicated coliform bacteria, and wells that were yellow and fluorescent when exposed to UV light (366 nm) indicated *E. coli*. Results were expressed as log CFU/25 g based upon the average from triplicate plates.

### **2.6.2. pH Analyses**

The pH was measured in a mixture of non-inoculated ground meat and distilled water (ratio 1:10) using a digital pH meter (Easy Five 720, Mettler Toledo, Switzerland) calibrated at three points (4, 7 and 10) [36]. Measurements were performed in triplicate.

### **2.7. Statistical Analyses**

The statistical differences between chemical parameters were determined by analysis of variance (one-way ANOVA) followed by a least significant difference test at a 95% confidence level.

### **3. RESULTS & DISCUSSION**

### 3.1. Wine Chemistry

The results of chemical analyses of the wines are shown in Table 1. The chemical parameters, alcohol, pH, and acidity in the wines were in the range of 12.8–14.3 vol%, 3.5–3.9, and 3.8–5.7 g/L respectively. Except for GM, the SO<sub>2</sub> was within the allowed limits (150 mg/L for red wines, 200 mg/L for white and rosé wines) [37]. Being a very complex process, winemaking involves steps that enrich the wine with components that enhance the flavor, color, and aroma. One such step, the maceration, has the purpose of extracting such compounds from the hard parts of grapes, which are left in contact with the grape juice during fermentation and later during the elaboration of the wine. For example, seven days of maceration resulted in a much higher anthocyanin content in red wine relative to the rosé obtained from the same grape. Similarly, the concentration of total phenols (TP) and tannins significantly increased with maceration, both in white and red wine (Table 1). The phenolic composition of the wine undergoes significant changes during the crushing of grapes, maceration, and fermentation [12,22,25].

In macerated Graševina and Plavac Mali, the concentration of total phenolics was previously reported to be 2699–2850 mg GAE/L and 3210 mg GAE/L, respectively, which is significantly higher in comparison to standard Graševina (305 mg GAE/L) and Plavac Mali Rosé (1074 mg GAE/L) [22].

Phenolic compounds participate in developing the sensory characteristics of wine, such as color, astringency, and bitterness, with most of these compounds having a wide range of biological properties including antimicrobial [9–11,20,25]. Some reports indicate that phenolics from wines are probably the most active components in inhibiting the growth of pathogens [10]. On the other hand, the synergy of different compounds such as ethanol, phenolics, certain organic acids, SO<sub>2</sub>, and low pH in reducing the bacterial count seems to remain the key to wine bioactivity [6,7,11,13].

### 3.2. Antimicrobial Activity

The results of the antimicrobial activity of wines are shown in Table 2. The MIC values show the minimal concentration of wine needed to kill *E. coli*, while the results of the agar-well diffusion method show the ability of the wine to inhibit the growth of inoculated bacteria. The wines demonstrated antimicrobial effect against both *E. coli* strains. The two wines with the lowest phenolic content (GS and PM 0) had higher antimicrobial activity in comparison with the other wines. Aside from the small concentration of total phenolics (Table 1), these two



wines had the highest content in total (124–173 mg/L) and free SO<sub>2</sub> (9–27 mg/L) previously pointed out as a possible inhibitor of bacterial growth in the grape juice and wine [13]. As the only wine with no added SO<sub>2</sub>, GM showed the lowest antimicrobial activity. Although studies showed a contribution of malic and tartaric acid in the inactivation of pathogens, particularly with the low surrounding pH ( $\leq 3$ ) [6,13], our results demonstrated that the wines with the same acidity (GS and GM) have significantly different MIC values (Table 2). On the other hand, PM 0 with the lowest acidity value (Table 1) showed good antimicrobial activity. It is important to note that phenolics alone are certainly not responsible for the antimicrobial effect of the wines. PM7 is the only wine that showed different MIC values for the two strains, with *E. coli* 12 being more sensitive; however, the mechanism behind this result was not determined. It can only be assumed that a particular group of compounds present in PM7 has more than one mechanism of action (discussed later in the text) on the bacterium resulting in a better antimicrobial effect.

Using the agar-disk and agar-well diffusion methods, we could not demonstrate antimicrobial effectiveness for any of the tested wines. Inhibition zones were not detected around the discs and around the wells they were <8 mm, which indicates no inhibitory effect [33]. The small volume of wine (25 and 50  $\mu$ L) and a correspondingly small amount of potentially antimicrobial wine components might be why the effect was not recorded. This brings to question the suitability of these methods for testing the antimicrobial activity of the wines. On the other hand, MIC values showed accurate, reproducible, and reliable results.

**Table 2.** Antimicrobial activity of wines ( $n = 6$ ).

Wine Samples *	MIC (mg GAE/L)		Well (mm)	
	<i>E. coli</i> 12	<i>E. coli</i> 13	<i>E. coli</i> 12	<i>E. coli</i> 13
GS	122.0	122.0	<8	<8
GM	1079.6	1079.6	<8	<8
GM + SO <sub>2</sub>	570.0	570.0	<8	<8
PM 0	215.6	215.6	<8	<8
PM 7	397.8	795.6	<8	<8
Ethanol (13–15%)	0	0	<8	<8

\* GS—standard white wine, GM and GM + SO<sub>2</sub>—macerated white wine with and without SO<sub>2</sub>, PM 0—rosé, PM 7— macerated red wine.

We could not establish the specific relationship between the winemaking process and a particular MIC value, taking into account the pH value, ethanol content, SO<sub>2</sub>, or total phenolics. The findings rather indicate the importance of their synergy for the observed overall

antimicrobial effect of the tested wines. The maceration duration and the addition of sulfur in the vinification process could not be fully correlated with the antimicrobial effects of the wines. This was also previously confirmed in several studies [6,12,14]. Further, the ethanol diluted to the level of concentration found in the wines (13 to 15%) showed no antimicrobial activity. Previous in vitro studies demonstrated that ethanol concentration in the wine (usually between 10 and 13% v/v) is considerably low to account for bactericidal effect [6,13,38]. The bactericidal impact of the wine itself, for a given ethanol concentration, is more effective than any other alcoholic beverage. Some authors find that the combination of organic acids (lactic, malic, tartaric) and ethanol contribute to this stronger antimicrobial effect of the wine [38–40].

The particular role of total phenolics remains unclear. In previous studies, it has been suggested that compounds such as flavonoids (quercetin and quercetin-3-glucoside) and monomeric anthocyanins might be used as biochemical markers that contribute to the antimicrobial activity of red wines [9,14]. Among wine phenolics, resveratrol has been particularly emphasized as a strong antimicrobial compound [34,41]. In line with our results, Boban et al. [42] showed that resveratrol levels in the intact, dealcoholized, and thermally treated wines at 125 °C, which were similar in pH, total phenolics and ethanol content, differed significantly, although the wines showed a similar antimicrobial effect. To clarify the role of polyphenols, pH, ethanol, and other wine components, a study tested the antimicrobial effects of intact wine in comparison to that of phenols-stripped wine, dealcoholized wine, ethanol, and low pH applied separately and in combination [23]. Antibacterial activity of the samples could not be related to their total phenolics and resveratrol content, ethanol content, or pH. After intact wine, the phenols-stripped wine had the strongest antimicrobial effect against *S. enterica* and *E. coli*, so the authors concluded that nonphenolic constituents of wine were responsible for a major part of its antimicrobial activity. Further studies are needed to clarify the mechanism of antimicrobial action of phenolic compounds and their correlation with other chemical parameters, but it can be concluded that the antimicrobial activity of a complex solution such as wine is based on more than one compound.

### 3.3. The Experiment on Fish Samples

The pH values of the wine-marinated fish meat used for the inoculation of bacteria are shown in Table 3.


**Table 3.** The pH of the wine-marinated fish meat ( $n = 3$ ).

Wine Samples *	pH
Water	6.58 ± 0.01 a
GS	6.27 ± 0.02 b
GM	6.32 ± 0.01 b
GM + SO <sub>2</sub>	6.40 ± 0.01 c
PM 0	6.25 ± 0.01 b
PM 7	6.28 ± 0.01 b

\* GS—standard white wine, GM and GM + SO<sub>2</sub>—macerated white wine with and without SO<sub>2</sub>, PM 0—rosé, PM 7— macerated red wine; <sup>a-c</sup> means followed by different superscript are significantly different ( $p < 0.05$ ).


In wine-marinated fish meat without inoculum, the number of CFU/25 g was zero in all samples. The total viable bacterial count or other bacteria genera were not analyzed in raw material or during the experiment.

Three days after the inoculation of the wine-marinated fish fillets with *E. coli* strains, which were held at  $4 \pm 1$  °C, the initial loads were  $<7$  log CFU/25 g in all samples (Figures 2 and 3). After three days of storage, the levels of both *E. coli* strains were the lowest in GM (biggest MIC values and no added sulfur in winemaking process) and GS samples, followed by the rosé and standard red wines (PM 0 and PM 7).



**Figure 2.** Count of *E. coli* (WDCM00012) in wine-marinated seabass (*Dicentrarchus labrax*); (GS—standard white wine, GM and GM + SO<sub>2</sub>—macerated white wine with and without SO<sub>2</sub>, PM 0—rosé, PM 7—macerated red wine) after 3 and 7 days of storage at 4 ± 1 °C.

The chemical parameters of wine samples could not be correlated with their antimicrobial activity. The 200 mL of wine that was used for marinating the fillets contained five times less TP than reported in Table 1, accordingly 61, 540, 570, 215, and 642 mg GAE in GS, GM, GM + SO<sub>2</sub>, PM 0, and PM 7, respectively. For GS, GM, and PM 0 the MIC values obtained are presented below, however, these wines showed better inhibition against *E. coli* strains than GM + SO<sub>2</sub> which contained the inhibitory concentration of TP in the 200 mL and high levels of SO<sub>2</sub>. PM 7, with the highest TP content (including anthocyanins and tannins), showed similar inhibition strength as other wines, with no difference between the two strains regardless of the MIC values. The pH with the contribution of ethanol was found to be an important factor in predicting in vitro inactivation and the efficacy of wine treatments against *E. coli* [6]. Our results show that this contribution is not valid for food model testing.



**Figure 3.** Count of *E. coli* (WDCM00013) in wine-marinated seabass (*Dicentrarchus labrax*); (GS—standard white wine, GM and GM + SO<sub>2</sub>—macerated white wine with and without SO<sub>2</sub>, PM 0—rosé, PM 7—macerated red wine) after 3 and 7 days of storage at 4 ± 1 °C.

Subsequent storage resulted in the growth of bacteria in all samples, from 0.6 to 1.5 log cycles in samples inoculated with *E. coli* 12, and from 0.6 to 1.4 log cycles in *E. coli* 13. After seven days of storage, the number of *E. coli* 12 was the lowest in GM and *E. coli* 13 in GS. The effect of rosé and red wines (PM 0 and PM 7) was similar, and the growth was again the highest in GM + SO<sub>2</sub>. In comparison to the control, all wines showed the capacity to reduce the number and growth of heavily infected sea bass filets. The role of SO<sub>2</sub> in the antimicrobial activity of wine is questionable; its purpose in wine seems to be mainly antioxidant (prevention of browning) [22].

The specific spoilage bacteria such as *Pseudomonas* and H<sub>2</sub>S-producing are mostly predominant in the spoilage flora of chilled seabass, causing off-flavors, and sensory rejection. Generally, the number of *Enterobacteriaceae* counts in chilled seabass is around 2 log CFU/g, and remain at low levels until ca. 7–9 days of storage [43]. The antimicrobial effect of phenolics against these microorganisms is usually concentration-dependent [44]. It has been reported that the phenolics in wine inhibit the growth of bacteria by multiple mechanisms, including membrane damage, inhibition of nucleic acid synthesis, inhibition of metabolism, and inhibition of cell wall and cell membrane synthesis [45]. The catechin was reported to retard the growth of spoilage bacteria in fish products [44], also high-tannin extracts were found to inhibit growth, in contrast to high-flavonoid extracts, which were found to induce bacterial growth [19]. There have been few studies conducted pertaining to bacterial survival, especially

*E. coli* in food models with wine. Boban et al. [42] found that the intact red wine kills *S. enterica* and *E. coli* after direct exposure in less than 5 min. In addition, the direct exposure of bacterial cells to white and red wine resulted in the rapid inactivation of *C. jejuni* [7]. The white and red wine marinades have shown beneficial effects against meat products contaminated with different foodborne pathogens [7,15,16,18]. In all of the studies mentioned, the wine was used as one of the ingredients of the marinades, and its antimicrobial effect was tested in combination with salt, spices, plant extracts, and essential oils. To our knowledge, there is no available literature on the challenge tests against *E. coli* using wine as the main antimicrobial substance. Being a source of the numerous bioactive compounds, it is hard to separate one distinctive segment responsible for wine's antimicrobial activity. The results showed sufficient evidence that the wines should be further investigated to relate the antimicrobial properties with its various compounds [38].

## **4. CONCLUSIONS**

This research aimed to evaluate the antimicrobial effect of different wines, produced using different technologies, against two strains of *E. coli*, and the strains' behavior in wine marinated seabass fillets. The study gives evidence that all wines have antimicrobial effects in vitro and in food models, and indicate that exposure to wine may inhibit the growth of the strains; however, the extrapolation of the results when the foods are contaminated with *E. coli* should be taken with reserve since a low number of strains were tested, invasiveness of the strains after treatment with wine was not determined, and the inhibitory effect was obtained during a short period of storage. Taking into account that in a complex nutrient medium such as fish, there are numerous reactive mechanisms between food and wine; thus the food matrix may have some buffering capacity which can reduce the antimicrobial effect. Further studies should involve other bacterial strains and genera, investigations of different microbial virulence factors, and antimicrobial effects of individual phenolic compounds and their mixtures to uncover mechanisms involved in the antimicrobial response.



## **5. REFERENCES**

1. Fung F, Wang H.S, Menon S. Food Safety in the 21st Century. *Biomed.* 2018;41:88–95.
2. Chen L, Alali W. Recent Discoveries in Human Serious Foodborne Pathogenic Bacteria: Resurgence, Pathogenesis, and Control Strategies. *Front. Microbiol.* 2018;9:2412.
3. Xu J.G, Cheng B.K, Jing H.Q. *Escherichia Coli* O157: H7 and Shiga-like-toxin-producing *Escherichia Coli* in China. *World J. Gastroenterol.* 1999;5:191–94.
4. Mylius S.D, Nauta M.J, Havelaar A.H. Cross-contamination during food preparation: A mechanistic model applied to chicken-borne *Campylobacter*. *Risk Anal. Int.* 2007;27:803–13.
5. van Elsas J.D, Semenov A.V, Costa R, Trevors J.T. Survival of *Escherichia coli* in the environment: Fundamental and public health aspects. ISME. 2011. doi:10.1038/ismej.2010.80.
6. Waite J.G, Daeschel M.A, Contribution of wine components to inactivation of food-borne pathogens. *Food Sci.* 2007;7:M286–M291.
7. Isohanni P, Alter T, Saris P, Lyhs U. Wines as possible meat marinade ingredients possess antimicrobial potential against *Campylobacter*. *Poult. Sci.* 2010;89:2704–10.
8. Kargiotou C, Katsanidis E, Rhoades J, Kontominas M, Koutsoumanis K. Efficacies of soy sauce and wine base marinades for controlling spoilage of raw beef. *Food Microbiol.* 2011;28:158–63.
9. Tenore G.C, Basile A, Novellino E. Antioxidant and antimicrobial properties of polyphenolic fractions from selected Moroccan red wines. *Food Sci.* 2011;76:C1342–C1348.
10. Papadopoulou C, Soulti K, Roussis I.G. Potential antimicrobial activity of red and white wine phenolic extracts against strains of *Staphylococcus aureus*, *Escherichia coli* and *Candida albicans*. *Food Technol. Biotechnol.* 2005;43:41–6.
11. Friedman M, Henika P.R, Levin C.E, Mandrell R.W. Antimicrobial wine formulations against the foodborne pathogens *Escherichia coli* O157:H7 and *Salmonella enterica*. *Food. Sci.* 2006;71:M245–M251.
12. Friedman M. Antibacterial, antiviral, and antifungal properties of wines and winery byproducts in relation to their flavonoid content. *Agric. Food Chem.* 2014;62:6025–42.
13. Moretro T, Daeschel M.A. Wine is bactericidal to food-borne pathogens. *Food Sci.* 2004;69:M250–M257.

14. Radovanović A.N, Jovančičević B.S, Radovanović B.C, Mihajilov-Krstev T. Antibimicrobial effectiveness of selected Vranac wines against six Gram-positive and six Gram-negative bacterial strains. *Trop. J. Pharm. Res.* 2014;13:819–24.
15. Birk T, Knöchel S. Fate of food-associated bacteria in pork as affected by marinade, temperature, and ultrasound. *Food Prot.* 2009;72:549–55.
16. Nisiotou A, Chorianopoulos N.G, Gounadaki A, Panagou E.Z, Nychas, G.J. Effect of wine-based marinades on the behavior of *Salmonella Typhimurium* and background flora in beef fillets. *Int. J. Food Microbiol.* 2013;164:119–27.
17. Rhoades J, Kargiotou C, Katsanidis E, Koutsoumanis K.P. Use of marination for controlling *Salmonella enterica* and *Listeria monocytogenes* in raw beef. *Food Microbiol.* 2013;36:248–53.
18. Díez J.G, Patarata L. Behavior of *Salmonella* spp., *Listeria monocytogenes*, and *Staphylococcus aureus* in Chourico de Vinho, a dry fermented sausage made from wine-marinated meat. *Food Prot.* 2013;76:588–94.
19. Friedman M, Levin C.E, Henika P.R. Addition of phytochemical-rich plant extracts mitigate the antimicrobial activity of essential oil/wine mixtures against *Escherichia coli* O157:H7 but not against *Salmonella enterica*. *Food Control.* 2017. doi:10.1016/j.foodcont.2016.09.002.
20. Rodríguez-Vaquero M.J, Aredes-Fernández P.A, Manca de Nadra M.C. Phenolic compounds from wine as natural preservatives of fish meat. *Food Technol. Biotechnol.* 2013;51:376–82.
21. Lee J.H, Cho B.K. Effects of grape, wild grape and black raspberry wines on ground pork during refrigerated storage. *Acta Aliment.* 2014;43:553–63.
22. Boban N, Tonkic M, Budimir D, Modun D, Sutlovic D, Punda-Polic V et al. Antimicrobial effects of wine: Separating the role of polyphenols, pH, ethanol, and other wine components. *Food Sci.* 2010;75:M322–M326.
23. Amerine M.A, Ough C.S. *Methods for Analysis of Musts and Wines.* 1st ed. New York: John Wiley and Sons; 1980. p. 181–85.
24. Katalinić V, Možina S.S, Skroza D, Generalić I, Abramović H, Miloš M et al. Polyphenolic profile, antioxidant properties and antimicrobial activity of grape skin extracts of 14 *Vitisvinifera* varieties grown in Dalmatia (Croatia). *Food Chem.* 2010;119: 715–23.
25. Ribéreau-Gayon P, Stonestreet E. Dosage des tanins du vin rouge et détermination de leur structure. *Chim. Anal.* 1966;48:188–96.

26. Ribéreau-Gayon P, Stonestreet E. Le dosage des anthocyanes dans le vin rouge. Bull. Soc. Chim. Fr. 1965;9:2649–52.
27. Skroza D, Šimat V, SmoleMožina S, Katalinić V, Boban N, Generalić Mekinić I. Interactions of resveratrol with other phenolics and activity against food-borne pathogens. Food Sci. Nutr. 2019. doi:10.1002/fsn3.1073.
28. Balouiri M, Sadiki M, Ibsoudaet S.K. Methods for in vitro evaluating antimicrobial activity: A review. Pharm. Anal. 2016;6:71–9.
29. Koohsari H, Ghaemi E.A, Shesh Poli M.S, Sadegh A. Evaluation of antibacterial activity of Lemon verbena (*Lippiacitriodora*) leaves. Ann. Biol. Res. 2013;4:52–5.
30. ISO. Microbiology of Food and Animal Feeding Stuffs—Horizontal Method for the Enumeration of Presumptive *E. coli*—Part 2: Colony Count Technique at 44 °C Using 5-Bromo-4-Chloro-3-Indolyl-Beta-D-Glucuronic Acid; ISO 16649–2; International Organization for Standardization: Geneva, Switzerland, 2001; p. 8.
31. Šimat V, Bogdanović T, Krželj M, Soldo A, Maršić-Lučić J. Differences in chemical, physical and sensory properties during shelf life assessment of wild and farmed gilthead sea bream (*Sparus aurata*, L.). J. Appl. Ichthyol. 2012. doi:10.1111/j.1439-0426.2011.01883.
32. L 149/1 Official Journal of the European Union. Regulation (EU) No 1308/2013 of the European Parliament and of the Council as Regards Wine-Growing Areas where the Alcoholic Strength may be Increased, Authorised Oenological Practices and Restrictions Applicable to the Production and Conservation of Grapevine Products, the Minimum Percentage of Alcohol for By-Products and Their Disposal, and Publication of OIV Files.
33. Milat A.M, Boban M, Teissedre P.-L, Šešelja-Perišin A, Jurić D, Skroza D et al. Effects of oxidation and browning of macerated white wine on its antioxidant and direct vasodilatory activity. Funct. Foods 2019;15:138–47.
34. Chan M.M. Antimicrobial effect of resveratrol on dermatophytes and bacterial pathogens of the skin. Biochem. Pharmacol. 2002;63: 99–104.
35. Boban N, Tonkić M, Modun D, Budimir D, Mudnić I, Sutlović D et al. Thermally treated wine retains antibacterial effects to food-borne pathogens. Food Control 2010;21:1161–65.
36. Cushnie T.T.P, Lamb A.J. Recent advances in understanding the antibacterial properties of flavonoids. Int. J. Antimicrob. Agents 2011;38:99–107.

37. CLSI. *Performance Standards for Antimicrobial Susceptibility Testing*; Twenty-Fourth Informational Supplement. CLSI document M100-S24; Clinical and Laboratory Standards Institute: Wayne, PA, USA, 2014.
38. CLSI. *Methods for Dilution Antimicrobial Susceptibility Tests for Bacteria That Grow Aerobically*; Approved Standard—8th ed. CLSI document M07-A8; Clinical and Laboratory Standards Institute: Wayne, PA, USA, 2009; p. 226.
39. The European Committee on Antimicrobial Susceptibility Testing. Breakpoint Tables for Interpretation of MICs and Zone Diameters. Version 10.0, 2020. Available online: <http://www.eucast.org>
40. Maqsood S, Benjakul S, Shahidi F. Emerging role of phenolic compounds as natural food additives in fish and fish products. *Food Sci. Nutr.* 2013. doi:10.1080/10408398.2010.518775.
41. Daglia M. Polyphenols as antimicrobial agents. *Curr. Opin. Biotechnol.* 2012. doi:10.1016/j.copbio.2011.08.007.
42. Papadopoulos V, Chouliara I, Badeka A, Savva I.N, Kontominas M.G. Effect of gutting on microbiological, chemical, and sensory properties of aquacultured sea bass (*Dicentrarchus labrax*) stored in ice. *Food Microbiol.* 2003. doi:10.1016/S0740-0020(02)00148-X.
43. Jamir L, Kumar V, Gat Y, Kumar A, Kaur S. Wine: A potential source of antimicrobial compounds. *Wine Res.* 2019. doi:10.1080/09571264.2019.1652151.
44. Just J.R, Daeschel M.A. Antimicrobial effects of wine on *Escherichia coli* O157: H7 and *Salmonella typhimurium* in a model stomach system. *Food Sci.* 2003;68:285–90.
45. Weisse M.E, Eberly B, Person D.A. Wine as a digestive aid: Comparative antimicrobial effects of bismuth salicylate and red and white wine. *BMJ.* 1995;311:1657–60.

## **6. SUMMARY**

**Objectives:** The main objective of the present study was to test the antimicrobial activity of five wines against two *E. coli* strains and to verify the findings in wine-marinated fish fillets as a food matrix.

**Materials and methods:** Five different wines (standard Graševina, macerated Graševina with and without sulfur, rosé, and standard Plavac Mali), all typical Croatian wines, were tested to determine the antimicrobial activity against two *Escherichia coli* bacterial strains (ATCC® 25922 and ATCC® 8739) in vitro and using sea bass (*Dicentrarchus labrax*) fillets as food matrix. The chemical composition of wines (pH, acidity, alcohol, total phenolics, anthocyanins, tannins, and sulfur content) and antimicrobial activity (minimal inhibitory concentration (MIC), agar-well diffusion method) were determined.

**Results:** The total phenolic content of the wines ranged from 305–3210 mg gallic acid equivalents per liter (GAE/L), and did not correlate to antimicrobial activity. The two wines with the lowest phenolic content (standard Graševina and rosé) had the lowest MIC values (122 and 429 mg GAE/L). A specific relation between the winemaking process and a particular MIC value was not established. There was also no relation found between the pH value, ethanol content, sulfur, or phenolics in regards to the antimicrobial effect. In fish fillets marinated in wine + water mixture ( $v/v = 1:1$ ) and inoculated with 7 log colony forming units (CFU)/25 g the growth of bacteria was reduced after three days of storage at 4 °C. Subsequent storage resulted in the growth of bacteria in all samples, with the lowest growth of *E. coli* ATCC® 25922 in macerated Graševina and *E. coli* ATCC® 8739 in standard Graševina. All wines showed the capacity to reduce the number and growth of heavily infected sea bass filets, but correlation with specific wine constituents was not found.

**Conclusions:** Taking into account the numerous reactive mechanisms between food and wine, all in vitro studies in controlled laboratory conditions should be further verified in the relevant environment, and additional research is needed to clarify the role of individual wine components in the mechanism of antimicrobial activity.

## **7. SAŽETAK**



**Ciljevi:** Glavni cilj istraživanja bio je dokazati antimikrobnu aktivnost pet vrsta vina na dva soja *Escherichiae coli* te potvrditi nalaze na prehrambenom proizvodu.

**Sredstva i metode:** Ispitana je *in vitro* antimikrobna aktivnost pet različitih vrsta vina (standardna Graševina, macerirana Graševina sa i bez sumpora, rosé i standardni Plavac Mali) na dva soja *Escherichia coli* (ATTC<sup>®</sup> 25922 i ATTC<sup>®</sup>8739) te u filetima brancina (*Dicentrachus labrax*) mariniranim u smjesi vina i vode (V/V=1:1) i inokuliranim bakterijama u koncentraciji 10<sup>7</sup> jedinica koje formiraju kolonije (CFU). Utvrđeni su slijedeći parametri vina: pH, kiselost, alkohol, ukupni fenoli, antocijani, tanini i udio sumpora te antimikrobna aktivnost korištenjem metode difuzije u jažicama i metode određivanja minimalne inhibitorne koncentracije (MIC).

**Rezultati:** Ukupan sadržaj fenola u vinima bio je u rasponu od 305 do 3210 mg izraženo u ekvivalentima galne kiseline po litri (GAE/L), a korelacija između sadržaja ukupnih fenola i antimikrobne aktivnosti nije utvrđena. Dvije vrste vina s najnižim udjelom ukupnih fenola (standardna Graševina i rosé) imali su i najniže MIC vrijednosti (122 i 429 mg GAE/L). Nije utvrđena povezanost tehnologije proizvodnje vina s dobivenim MIC vrijednostima. Također, nije utvrđena povezanost između pH vrijednosti, udjela etanola ili sumpora i antimikrobne aktivnosti ispitivanih vina. U mariniranim ribljim filetima, rast bakterija bio je smanjen nakon tri dana pohrane na 4 °C. Daljnja pohrana rezultirala je porastom bakterija u svim uzorcima, a najniži porast *E. coli* ATTC<sup>®</sup> 25922 zabilježen je u maceriranoj Graševini i *E. coli* ATTC<sup>®</sup>8739 u standardnoj Graševini. Sva vina pokazala su djelotvornost u inhibiciji rasta bakterija u filetima brancina inokuliranih velikim brojem CFU. Povezanost s pojedinačnim komponentama vina nije utvrđena.

**Zaključak:** Zbog mogućih brojnih ineterrekcija između kompleksnih medij kao što su prehrambeni proizvodi i vino, sva *in vitro* mikrobiološka istraživanja u kontroliranim laboratorijskim uvjetima trebala bi se potvrditi i u relevantnom okruženju. Potrebna su dodatna istraživanja kako bi se pojasnila uloga pojedinih komponenti vina u njegovom antimikrobnom djelovanju.

## **8. CURRICULUM VITAE**

HEIDI-CHRISTINE SANTORO

Hektorovićeva 19, 21000 Split Croatia / 0994223121 / [heidi.christine.santoro@gmail.com](mailto:heidi.christine.santoro@gmail.com)

Date of birth: 21/6/1987

WORK EXPERIENCE

2012 – Present

OWNER/CEO

Owner and CEO of heidichristine LLC, a multifaceted writing firm

EDUCATION

2003 – 2005

PPL & CPL – Scottsdale Private Airports

2004 – 2008

Undergraduate Degree in Linguistics and Communication – Obtained at: Arizona State University, Höskolan i Kalmar, Sweden, and St. Johns University, Rome

2009 – 2012

Columbia Law

2012 – 2013

National Paramedic Certification (NPR), Scottsdale Community College

EXTRA-CURRICULAR ACTIVITES

1996 – 2002

Competitive gymnast for South Africa and the USA

2012 – Present



Dancer/acrobat

HOBBIES

Sailing – I currently hold the boat skipper license category B

Article

# Antimicrobial Activity of Selected Red and White Wines against *Escherichia coli*: In Vitro Inhibition Using Fish as Food Matrix

Heidi Christine Santoro <sup>1</sup>, Danijela Skroza <sup>2,\*</sup> , Anđela Dugandžić <sup>2</sup>, Mladen Boban <sup>1</sup> and Vida Šimat <sup>3,\*</sup> 

<sup>1</sup> Department of Pharmacology, School of Medicine, University of Split, Šoltanska 2, HR-21000 Split, Croatia; heidi.christine.santoro@gmail.com (H.C.S.); mladen.boban@mefst.hr (M.B.)

<sup>2</sup> Department of Food Technology and Biotechnology, Faculty of Chemistry and Technology, University of Split, Ruđera Boškovića 35, HR-21000 Split, Croatia; angy.duga96@gmail.com

<sup>3</sup> Department of Marine Studies, University of Split, Ruđera Boškovića 37, HR-21000 Split, Croatia

\* Correspondence: danci@ktf-split.hr (D.S.); vida@unist.hr (V.Š.); Tel.: +385-21-329-458 (D.S.); +385-21-510-192 (V.Š.)

Received: 15 June 2020; Accepted: 14 July 2020; Published: 15 July 2020



**Abstract:** Five different wines (standard Graševina, macerated Graševina with and without sulfur, rosé, and standard Plavac Mali), all typical Croatian wines, were tested to determine the antimicrobial activity against two *Escherichia coli* bacterial strains (ATCC<sup>®</sup> 25922 and ATCC<sup>®</sup> 8739) in vitro and using sea bass (*Dicentrarchus labrax*) fillets as food matrix. The chemical composition of wines (pH, acidity, alcohol, total phenolics, anthocyanins, tannins, and sulfur content) and antimicrobial activity (minimal inhibitory concentration (MIC), agar-well diffusion method) were determined. The total phenolic content of the wines ranged from 305–3210 mg gallic acid equivalents per liter (GAE/L), and did not correlate to antimicrobial activity. The two wines with the lowest phenolic content (standard Graševina and rosé) had the lowest MIC values (122 and 429 mg GAE/L). A specific relation between the winemaking process and a particular MIC value was not established. There was also no relation found between the pH value, ethanol content, sulfur, or phenolics in regards to the antimicrobial effect. In fish fillets marinated in wine + water mixture (*v/v* = 1:1) and inoculated with 7 log colony forming units (CFU)/25 g the growth of bacteria was reduced after three days of storage at 4 °C. Subsequent storage resulted in the growth of bacteria in all samples, with the lowest growth of *E. coli* ATCC<sup>®</sup> 25922 in macerated Graševina and *E. coli* ATCC<sup>®</sup> 8739 in standard Graševina. All wines showed the capacity to reduce the number and growth of heavily infected sea bass filets, but correlation with specific wine constituents was not found. Taking into account the numerous reactive mechanisms between food and wine, all in vitro studies in controlled laboratory conditions should be further verified in the relevant environment, and additional research is needed to clarify the role of individual wine components in the mechanism of antimicrobial activity.

**Keywords:** wine; sea bass; antimicrobial activity; MIC; phenols

## 1. Introduction

Food-borne bacteria have become an ever-increasing challenge, in both commercial and domestic use. Contaminated food continues to cause millions of people to fall ill, with many succumbing to fatality due to foodborne bacterial infections [1]. *Escherichia coli* are among the most common food-borne pathogens related to food spoilage and food poisoning. *E. coli* are found in the environment and the lower intestines of warm-blooded organisms as a part of normal microflora and represent mostly harmless bacteria. Certain types of *E. coli*, such as shiga-like-toxin-producing *E. coli* (serotype

O157:H7, SLTEC), contaminate different types of foods and water, presenting a risk for humans [2,3]. The main sources of human infections are undercooked and contaminated meat and fish, as well as dairy products made from raw milk. Despite various measures taken during processing, consumers can still be exposed to this pathogen by consumption and handling of improperly prepared foods or through cross-contamination as another major risk factor [3,4].

*E. coli* can tolerate and rapidly adapt to a variety of stressful conditions such as fluctuations and extremes of temperature, high osmolarity, and low pH [5]. One potential approach to reducing the counts of *E. coli* could be the application of natural and sustainable substances with antimicrobial properties. Red and white wines have demonstrated antimicrobial activity under various experimental conditions against spoilage and pathogenic food bacteria such as *Bacillus* spp., *Escherichia* spp., *Listeria* spp., *Salmonella* spp., *Sigella*, *Pseudomonas* spp., *Campylobacter jejuni*, *Yersinia enterocolitica*, *Vibrio* spp., and *Staphylococcus aureus* [6–19].

It is generally held that wine's biological potential is related to a large number of polyphenolic compounds [12]. Polyphenols (phenolics) are plant secondary metabolites, which have essential roles in plant physiology but have potential health benefits on the human organism, mainly as antioxidant, anti-allergic, anti-inflammatory, anticancer, antihypertensive, and antimicrobial agents [20]. The antimicrobial properties of polyphenols have been widely studied to propose new methods of food preservation and enhance food safety without the use of synthetic additives. Among polyphenol classes, *E. coli* showed sensitivity to phenolic acids, hydrolysable and condensed tannins, flavan-3-ols, and flavonols. All of these are found in fruits and vegetables, and plant-derived beverages, such as tea and wine [20]. Red and white wines contain a variety of phenolic compounds that have been associated with numerous health benefits mentioned earlier but have also been active against foodborne pathogens.

In past years, great attention has been paid to wine polyphenols and naturally produced macerated white wines are becoming increasingly popular; however their biological effects are rarely studied. These wines have a high content of biologically active compounds and among them; polyphenols are considered the main factor responsible for particular biological properties. The phenolic composition of wines varies with a grape variety, but it is also influenced by viticultural and environmental factors, and the wine-making process. However, the skins and seeds of grapes are known to be rich sources of polyphenolic compounds, both flavonoids and non-flavonoids [21], and thus phenolic-rich wines are commonly obtained through maceration step, in which grape juice is left in contact with grape skins, seeds and even stems, for weeks and even months [22]. Although wine polyphenols have been intensively studied, it has been demonstrated that the synergistic effect of phenolics with other wine components, such as low pH, ethanol, and organic acids, remains the key mechanism of wine bioactivity [11,12,23]. The specific components of wine and their relative contribution to its antimicrobial activity are still a matter of debate.

There have been a limited number of studies investigating the antimicrobial effect of wine used in the process of food preparation, either as an ingredient or marinade. In available studies, the wine was not tested intact, but in the form of wine extract or in the combination with soya sauce, oregano and thyme essential oils, garlic juice, or even pure phenolic compounds [7–10,24,25]. Marinade recipes containing red or white wine enriched with plant essential oils demonstrated strong activity under laboratory conditions against major foodborne pathogens [11]. Also, the wine application in foods contributed to the prolongation of shelf life of meat [8] and fish [24] products with the ability to reduce total viable count in the products. Both red and white wines from different regions and grape varieties showed the antimicrobial potential against *E. coli*, however, the in vitro effect was not often confirmed in food models [12,19].

There is an increasing interest in investigations into the use of bioactive natural compounds for the enhancement of food safety and product quality. Microbiological challenge tests are an important tool necessary for simulation of assessment of real-life food safety and prediction of safety risks. The use of natural antimicrobials and preservative sources in foods such as wine would help reduce the use of synthetic additives and aggressive processing, without compromising food safety. In this regard, the main objective of the present study was to test the antimicrobial activity of five wines against two *E. coli* strains and to verify the findings in wine-marinated fish fillets as a food matrix.

## 2. Materials and Methods

### 2.1. Wine Samples and Technology

The five wines, three white wines produced from the grape variety Graševina in 2015 at the Krauthaker winery, and one red and one rosé produced from the grape variety Plavac Mali in 2015 at the Volarević winery, Croatia, were tested *in vitro* for antimicrobial activity in the standard growing media and in fish fillets as food matrix. The technology of production of the wines tested is schematically shown in Figure 1. The standard white wine (GS) and rosé wine (PM 0) were produced following a classic methodology requiring grape juice to be separated from the hard parts of the grapes during fermentation, without maceration. Conversely, the Georgian wine production principles were used in obtaining macerated white wines rich in polyphenols. Following spontaneous fermentation, and without removing grape seeds and skins, the tanks were sealed air-tight, allowing the wine further elaboration for four months. Part of the wine so produced was bottled without further treatment (GM), while sulphur (GM + SO<sub>2</sub>) was added to the other part as a preservative. The red wine (PM 7) production process included both fermentation and maceration for seven days in contact with the grape seeds and skins (Figure 1).




Figure 1. Technology of the tested wines.

### 2.2. Wine Chemistry Analyses

The enological analysis of all wines included ethanol and sulfur content, along with pH and acidity values. These were determined according to the Compendium of International Methods of Analysis of Wines and Musts of the International Organisation of Vine and Wine (OIV), and are shown in Table 1.

**Table 1.** Chemical composition of tested wines.

Wine Samples *	pH	Acidity (g/L) **	Alcohol (%)	Total Phenolics (mg GAE/L)	Anthocyanins (mg/L)	Tannins (mg/L)	Free SO <sub>2</sub> (mg/L)	Total SO <sub>2</sub> (mg/L)
GS	3.54	4.9	13.0	305 ± 3.5 <sup>a</sup>	0 <sup>a</sup>	0.26 ± 0.01 <sup>a</sup>	27	124
GM	3.94	4.8	13.3	2699 ± 8.2 <sup>b</sup>	0 <sup>b</sup>	3.34 ± 0.10 <sup>b</sup>	1	3
GM + SO <sub>2</sub>	3.90	5.4	12.8	2850 ± 34.6 <sup>b</sup>	0 <sup>c</sup>	4.93 ± 0.19 <sup>c</sup>	5	99
PM 0	3.54	3.8	14.3	539 ± 22.2 <sup>c</sup>	1.97 ± 0.04 <sup>b</sup>	1.08 ± 0.07 <sup>d</sup>	9	173
PM 7	3.61	5.7	13.7	1988.96 ± 35.0 <sup>d</sup>	30.14 ± 6.52 <sup>d</sup>	4.93 ± 0.46 <sup>e</sup>	6	89

\* GS—standard white wine, GM and GM + SO<sub>2</sub>—macerated white wine with and without SO<sub>2</sub>, PM 0—rosé, PM 7—macerated red wine; \*\* expressed as tartaric acid; <sup>a–e</sup> mean value ± standard deviation in the same column followed by different superscript are significantly different ( $p < 0.05$ ).

### 2.3. Analysis of Total Phenolics, Tannins and Anthocyanins

The total phenolic content of the wine samples was measured spectrophotometrically using the Folin-Ciocalteu method [21,26] with the results expressed as mg of gallic acid equivalents per liter (mg GAE/L).

The total tannin contents were measured by acidic hydrolysis of proanthocyanidins resulting in carbocation formation with partial conversion into red cyanidin using the method of Ribéreau-Gayon and Stonestreet [27] and concentration was obtained in g/L.

Anthocyanin content was determined by the SO<sub>2</sub> bleaching procedure and results are expressed as mg/L [28].

All measurements were carried out in triplicate using a SPECORD 200 Plus (Analytik Jena AG, Jena, Germany).

### 2.4. The *Escherichia coli* Strains and Inoculum Preparation

Bacterial strains *E. coli* WDCM00012 (ATCC8739, *E. coli* 12) and WDCM00013 (ATCC25922, *E. coli* 13) were used as test microorganisms. Determination of antimicrobial activity was performed using Mueller Hinton Broth/Agar (MHB, MHA; Biolife, Italy). The cell culture suspension was prepared by selection of a colony from the 24-hr old MHA plates, with adjustment in MHB to include a concentration of approximately 10<sup>8</sup> colony forming units per mL (CFU/mL) according to the turbidity of 0.5 McFarland scale standard (Densomat, bioMerieux, Marcy l'Etoile, France). An aliquot of 2 mL of cell culture suspension was diluted with 38 mL of MHB medium for inoculum preparation (initial concentration 10<sup>7</sup> CFU/mL).

### 2.5. In Vitro Evaluation of Antimicrobial Activity of Wines

#### 2.5.1. Agar-Disk and Agar-Well Diffusion Assays

The protocols used for agar-disk and agar-well diffusion assays are defined by Clinical and Laboratory Standards Institute (CLSI) [29,30]. The Petri plates containing 20 mL MHA were inoculated by spreading microbial inoculum over the entire agar surface using sterile cotton swabs rolled in the inoculum. For the disk diffusion method, filter paper discs of 6 mm in diameter containing 25 µL of the wine samples were placed on the agar surface. The same was done for the antibiotics, chloramphenicol, gentamicin, and tigecycline, used as control. The Petri plates were incubated at 37 °C for 20–24 h. The inhibition zone of the antibiotics was in the range proposed by CLSI and the European Committee on Antimicrobial Susceptibility Testing (EUCAST) [30,31] for both *E. coli* strains.

For agar-well diffusion assays, four equidistant holes were made in the agar using sterile cork borers (Ø = 7 mm) and 50 µL of the wine samples was added directly in the wells. The agar plates were at 4 °C for one hour, following incubation at 37 °C for 20–24 h. The inhibition zones formed on the medium were measured in millimeters (mm) [32]. If the inhibition zone was ≥12 mm, the tested sample was considered to have good inhibitory effect [33]. All measurements were done in triplicate with two positive growth controls included (ethanol in the concentration of 13–15%).

### 2.5.2. Minimal Inhibitory Concentration (MIC)

Minimal inhibitory concentration was measured according to the previously described method by [32,34]. Shortly 50  $\mu$ L of bacterial inoculum and 50  $\mu$ L of two-fold serially diluted wine samples were added to the wells of a sterile 96-well microtitre plate. Three antibiotics (chloramphenicol, gentamicin, and tigecycline) and ethanol (13–15% vol) were used as control. The MIC values of chloramphenicol, gentamicin, and tigecycline were in the range proposed by CLSI and EUCAST standards, 4, 0.5 and 0.15  $\mu$ g/mL, respectively [30,31]. The MIC was defined as the lowest concentration displaying the ability to inhibit any visible bacterial growth after adding 10  $\mu$ L/well of INT (2-p-iodophenyl-3-p-nitrophenyl-5-phenyl tetrazolium chloride, Sigma, Darmstadt, Germany). All measurements of MIC values were done in triplicate and expressed as mg GAE/L.

### 2.6. Preparation of Fish Samples and Inoculation with *E. coli*

Seabass (*Dicentrarchus labrax*) were purchased at the local fish market and transported to the lab packed with ice in polystyrene boxes. The fish were of class L, weighing ~400–600 g each. All further preparations were performed under sterile conditions. The fish was filleted and the fillets thoroughly washed in sterile distilled water. Five groups of fish fillets, each weighing approximately 200 g, were marinated in a mixture of wine and sterile distilled water (400 mL) prepared in the ratio 1:1. The control sample was kept in sterile distilled water. After two hours, each group of fillets was drained and ground in a knife mill down to 40–50 mm particles (GRINDOMIX GM 200, Retsch GmbH, Haan, Germany). Next, 50 g of ground meat was separated for pH measurement.

For the challenge test experiment, triplicate packages containing chunks of  $25 \pm 0.01$  g of ground wine-marinated fish meat were placed in sterile vacuum bags, inoculated with *E. coli* strains (0.5 mL of liquid inoculum prepared as described in Section 2.4) to achieve an initial value of  $\sim 7$  log CFU/25 g and vacuum sealed. The samples were kept at room temperature for 10 min to allow possible attachment and diffusion [7] and then stored at  $4 \pm 1$  °C. The number of *E. coli* was enumerated after three and seven days. In addition, triplicate packages of wine-marinated fish meat without inoculum were tested for *E. coli*.

#### 2.6.1. Enumeration of *E. coli*

The enumeration of *E. coli* was done using ISO methodology [35]. Briefly, the samples were transferred to stomacher bags with 225 mL of sterile peptone water and homogenized using a laboratory blender (Stomacher, Maxicator, IUL S.A., Barcelona, Spain) for three minutes. The serial of decimal dilutions was prepared and distributed to sterile Petri dish followed by the Tryptone Bile X-GLUC agar (Biolife, Milan, Italy) and incubated at 44 °C for 24–48 h. Plates with 10 to 150 blue-green colonies were counted. Confirmation of *E. coli* was done by Colilert-18 (Quanti-Tray/2000) method. According to the manufacturer's instructions, yellow wells indicated coliform bacteria, and wells that were yellow and fluorescent when exposed to UV light (366 nm) indicated *E. coli*. Results were expressed as log CFU/25 g based upon the average from triplicate plates.

#### 2.6.2. pH Analyses

The pH was measured in a mixture of non-inoculated ground meat and distilled water (ratio 1:10) using a digital pH meter (Easy Five 720, Mettler Toledo, Switzerland) calibrated at three points (4, 7 and 10) [36]. Measurements were performed in triplicate.

### 2.7. Statistical Analyses

The statistical differences between chemical parameters were determined by analysis of variance (one-way ANOVA) followed by a least significant difference test at a 95% confidence level.



### 3. Results and Discussion

#### 3.1. Wine Chemistry

The results of chemical analyses of the wines are shown in Table 1. The chemical parameters, alcohol, pH, and acidity in the wines were in the range of 12.8–14.3 vol%, 3.5–3.9, and 3.8–5.7 g/L respectively. Except for GM, the SO<sub>2</sub> was within the allowed limits (150 mg/L for red wines, 200 mg/L for white and rosé wines) [37]. Being a very complex process, winemaking involves steps that enrich the wine with components that enhance the flavor, color, and aroma. One such step, the maceration, has the purpose of extracting such compounds from the hard parts of grapes, which are left in contact with the grape juice during fermentation and later during the elaboration of the wine. For example, seven days of maceration resulted in a much higher anthocyanin content in red wine relative to the rosé obtained from the same grape. Similarly, the concentration of total phenols (TP) and tannins significantly increased with maceration, both in white and red wine (Table 1). The phenolic composition of the wine undergoes significant changes during the crushing of grapes, maceration, and fermentation [12,22,25].

In macerated Graševina and Plavac Mali, the concentration of total phenolics was previously reported to be 2699–2850 mg GAE/L and 3210 mg GAE/L, respectively, which is significantly higher in comparison to standard Graševina (305 mg GAE/L) and Plavac Mali Rosé (1074 mg GAE/L) [22].

Phenolic compounds participate in developing the sensory characteristics of wine, such as color, astringency, and bitterness, with most of these compounds having a wide range of biological properties including antimicrobial [9–11,20,25]. Some reports indicate that phenolics from wines are probably the most active components in inhibiting the growth of pathogens [10]. On the other hand, the synergy of different compounds such as ethanol, phenolics, certain organic acids, SO<sub>2</sub>, and low pH in reducing the bacterial count seems to remain the key to wine bioactivity [6,7,11,13].

#### 3.2. Antimicrobial Activity

The results of the antimicrobial activity of wines are shown in Table 2. The MIC values show the minimal concentration of wine needed to kill *E. coli*, while the results of the agar-well diffusion method show the ability of the wine to inhibit the growth of inoculated bacteria. The wines demonstrated antimicrobial effect against both *E. coli* strains. The two wines with the lowest phenolic content (GS and PM 0) had higher antimicrobial activity in comparison with the other wines. Aside from the small concentration of total phenolics (Table 1), these two wines had the highest content in total (124–173 mg/L) and free SO<sub>2</sub> (9–27 mg/L) previously pointed out as a possible inhibitor of bacterial growth in the grape juice and wine [13]. As the only wine with no added SO<sub>2</sub>, GM showed the lowest antimicrobial activity. Although studies showed a contribution of malic and tartaric acid in the inactivation of pathogens, particularly with the low surrounding pH ( $\leq 3$ ) [6,13], our results demonstrated that the wines with the same acidity (GS and GM) have significantly different MIC values (Table 2). On the other hand, PM 0 with the lowest acidity value (Table 1) showed good antimicrobial activity. It is important to note that phenolics alone are certainly not responsible for the antimicrobial effect of the wines. PM7 is the only wine that showed different MIC values for the two strains, with *E. coli* 12 being more sensitive; however, the mechanism behind this result was not determined. It can only be assumed that a particular group of compounds present in PM7 has more than one mechanism of action (discussed later in the text) on the bacterium resulting in a better antimicrobial effect.

Using the agar-disk and agar-well diffusion methods, we could not demonstrate antimicrobial effectiveness for any of the tested wines. Inhibition zones were not detected around the discs and around the wells they were <8 mm, which indicates no inhibitory effect [33]. The small volume of wine (25 and 50  $\mu$ L) and a correspondingly small amount of potentially antimicrobial wine components might be why the effect was not recorded. This brings to question the suitability of these methods for testing the antimicrobial activity of the wines. On the other hand, MIC values showed accurate, reproducible, and reliable results.

**Table 2.** Antimicrobial activity of wines ( $n = 6$ ).

Wine Samples *	MIC (mg GAE/L)		Well (mm)	
	<i>E. coli</i> 12	<i>E. coli</i> 13	<i>E. coli</i> 12	<i>E. coli</i> 13
GS	122.0	122.0	<8	<8
GM	1079.6	1079.6	<8	<8
GM + SO <sub>2</sub>	570.0	570.0	<8	<8
PM 0	215.6	215.6	<8	<8
PM 7	397.8	795.6	<8	<8
Ethanol (13–15%)	0	0	<8	<8

\* GS—standard white wine, GM and GM + SO<sub>2</sub>—macetared white wine with and without SO<sub>2</sub>, PM 0—rosé, PM 7—macerated redwine.

We could not establish the specific relationship between the winemaking process and a particular MIC value, taking into account the pH value, ethanol content, SO<sub>2</sub>, or total phenolics. The findings rather indicate the importance of their synergy for the observed overall antimicrobial effect of the tested wines. The maceration duration and the addition of sulfur in the vinification process could not be fully correlated with the antimicrobial effects of the wines. This was also previously confirmed in several studies [6,12,14]. Further, the ethanol diluted to the level of concentration found in the wines (13 to 15%) showed no antimicrobial activity. Previous in vitro studies demonstrated that ethanol concentration in the wine (usually between 10 and 13% *v/v*) is considerably low to account for bactericidal effect [6,13,38]. The bactericidal impact of the wine itself, for a given ethanol concentration, is more effective than any other alcoholic beverage. Some authors find that the combination of organic acids (lactic, malic, acidic, and tartaric) and ethanol contribute to this stronger antimicrobial effect of the wine [38–40].

The particular role of total phenolics remains unclear. In previous studies, it has been suggested that compounds such as flavonoids (quercetin and quercetin-3-glucoside) and monomeric anthocyanins might be used as biochemical markers that contribute to the antimicrobial activity of red wines [9,14]. Among wine phenolics, resveratrol has been particularly emphasized as a strong antimicrobial compound [34,41]. In line with our results, Boban et al. [42] showed that resveratrol levels in the intact, dealcoholized, and thermally treated wines at 125 °C, which were similar in pH, total phenolics and ethanol content, differed significantly, although the wines showed a similar antimicrobial effect. To clarify the role of polyphenols, pH, ethanol, and other wine components, a study tested the antimicrobial effects of intact wine in comparison to that of phenols-stripped wine, dealcoholized wine, ethanol, and low pH applied separately and in combination [23]. Antibacterial activity of the samples could not be related to their total phenolics and resveratrol content, ethanol content, or pH. After intact wine, the phenols-stripped wine had the strongest antimicrobial effect against *S. enterica* and *E. coli*, so the authors concluded that nonphenolic constituents of wine were responsible for a major part of its antimicrobial activity. Further studies are needed to clarify the mechanism of antimicrobial action of phenolic compounds and their correlation with other chemical parameters, but it can be concluded that the antimicrobial activity of a complex solution such as wine is based on more than one compound.

### 3.3. The Experiment on Fish Samples

The pH values of the wine-marinated fish meat used for the inoculation of bacteria are shown in Table 3.


**Table 3.** The pH of the wine-marinated fish meat ( $n = 3$ ).

Wine Samples *	pH
Water	6.58 ± 0.01 <sup>a</sup>
GS	6.27 ± 0.02 <sup>b</sup>
GM	6.32 ± 0.01 <sup>b</sup>
GM + SO <sub>2</sub>	6.40 ± 0.01 <sup>c</sup>
PM 0	6.25 ± 0.01 <sup>b</sup>
PM 7	6.28 ± 0.01 <sup>b</sup>

\* GS—standard white wine, GM and GM + SO<sub>2</sub>—macerated white wine with and without SO<sub>2</sub>, PM 0—rosé, PM 7—macerated red wine; <sup>a-c</sup> means followed by different superscript are significantly different ( $p < 0.05$ ).


In wine-marinated fish meat without inoculum, the number of CFU/25 g was zero in all samples. The total viable bacterial count or other bacteria genera were not analyzed in raw material or during the experiment.

Three days after the inoculation of the wine-marinated fish fillets with *E. coli* strains, which were held at  $4 \pm 1$  °C, the initial loads were  $<7$  log CFU/25 g in all samples (Figures 2 and 3). After three days of storage, the levels of both *E. coli* strains were the lowest in GM (biggest MIC values and no added sulfur in winemaking process) and GS samples, followed by the rosé and standard red wines (PM 0 and PM 7).



**Figure 2.** Count of *E. coli* (WDCM00012) in wine-marinated seabass (*Dicentrarchus labrax*); (GS—standard white wine, GM and GM + SO<sub>2</sub>—macerated white wine with and without SO<sub>2</sub>, PM 0—rosé, PM 7—macerated red wine) after 3 and 7 days of storage at  $4 \pm 1$  °C.

The chemical parameters of wine samples could not be correlated with their antimicrobial activity. The 200 mL of wine that was used for marinating the fillets contained five times less TP than reported in Table 1, accordingly 61, 540, 570, 215, and 642 mg GAE in GS, GM, GM + SO<sub>2</sub>, PM 0, and PM 7, respectively. For GS, GM, and PM 0 the MIC values obtained are presented below, however, these wines showed better inhibition against *E. coli* strains than GM + SO<sub>2</sub> which contained the inhibitory concentration of TP in the 200 mL and high levels of SO<sub>2</sub>. PM 7, with the highest TP content (including anthocyanins and tannins), showed similar inhibition strength as other wines, with no difference between the two strains regardless of the MIC values. The pH with the contribution of ethanol was found to be an important factor in predicting in vitro inactivation and the efficacy of wine treatments against *E. coli* [6]. Our results show that this contribution is not valid for food model testing.



**Figure 3.** Count of *E. coli* (WDCM00013) in wine-marinated seabass (*Dicentrarchus labrax*); (GS—standard white wine, GM and GM + SO<sub>2</sub>—macetared white wine with and without SO<sub>2</sub>, PM 0—rosé, PM 7—macerated redwine) after 3 and 7 days of storage at 4 ± 1 °C.

Subsequent storage resulted in the growth of bacteria in all samples, from 0.6 to 1.5 log cycles in samples inoculated with *E. coli* 12, and from 0.6 to 1.4 log cycles in *E. coli* 13. After seven days of storage, the number of *E. coli* 12 was the lowest in GM and *E. coli* 13 in GS. The effect of rosé and red wines (PM 0 and PM 7) was similar, and the growth was again the highest in GM + SO<sub>2</sub>. In comparison to the control, all wines showed the capacity to reduce the number and growth of heavily infected sea bass filets. The role of SO<sub>2</sub> in the antimicrobial activity of wine is questionable; its purpose in wine seems to be mainly antioxidant (prevention of browning) [22].

The specific spoilage bacteria such as *Pseudomonas* and H<sub>2</sub>S-producing are mostly predominant in the spoilage flora of chilled seabass, causing off-flavors, and sensory rejection. Generally, the number of *Enterobacteriaceae* counts in chilled seabass is around 2 log CFU/g, and remain at low levels until ca. 7–9 days of storage [43]. The antimicrobial effect of phenolics against these microorganisms is usually concentration-dependent [44]. It has been reported that the phenolics in wine inhibit the growth of bacteria by multiple mechanisms, including membrane damage, inhibition of nucleic acid synthesis, inhibition of metabolism, and inhibition of cell wall and cell membrane synthesis [45]. The catechin was reported to retard the growth of spoilage bacteria in fish products [44], also high-tannin extracts were found to inhibit growth, in contrast to high-flavonoid extracts, which were found to induce bacterial growth [19]. There have been few studies conducted pertaining to bacterial survival, especially *E. coli* in food models with wine. Boban et al. [42] found that the intact red wine kills *S. enterica* and *E. coli* after direct exposure in less than 5 min. In addition, the direct exposure of bacterial cells to white and red wine resulted in the rapid inactivation of *C. jejuni* [7]. The white and red wine marinades have shown beneficial effects against meat products contaminated with different foodborne pathogens [7,15,16,18]. In all of the studies mentioned, the wine was used as one of the ingredients of the marinades, and its antimicrobial effect was tested in combination with salt, spices, plant extracts, and essential oils. To our knowledge, there is no available literature on the challenge tests against *E. coli* using wine as the main antimicrobial substance. Being a source of the numerous bioactive compounds, it is hard to separate one distinctive segment responsible for wine's antimicrobial activity. The results showed sufficient evidence that the wines should be further investigated to relate the antimicrobial properties with its various compounds [38].

#### 4. Conclusions

This research aimed to evaluate the antimicrobial effect of different wines, produced using different technologies, against two strains of *E. coli*, and the strains' behavior in wine marinated seabass fillets. The study gives evidence that all wines have antimicrobial effects in vitro and in food models, and indicate that exposure to wine may inhibit the growth of the strains; however, the extrapolation of the results when the foods are contaminated with *E. coli* should be taken with reserve since a low number of strains were tested, invasiveness of the strains after treatment with wine was not determined, and the inhibitory effect was obtained during a short period of storage. Taking into account that in a complex nutrient medium such as fish, there are numerous reactive mechanisms between food and wine; thus the food matrix may have some buffering capacity which can reduce the antimicrobial effect. Further studies should involve other bacterial strains and genera, investigations of different microbial virulence factors, and antimicrobial effects of individual phenolic compounds and their mixtures to uncover mechanisms involved in the antimicrobial response.

**Author Contributions:** Conceptualization and methodology, D.S. and V.Š.; formal analysis and investigation D.S., V.Š., H.C.S., and A.D.; data curation, D.S., A.D. and V.Š.; writing—original draft preparation, D.S. and V.Š.; writing—review and editing, D.S., V.Š., H.C.S. and M.B.; visualization, D.S. and V.Š.; supervision, M.B. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research was funded by the Croatian Science Foundation (Project No. 8652).

**Acknowledgments:** Authors would like to thank the company Centaurus d.o.o., Split, Croatia for help with the fish material.

**Conflicts of Interest:** The authors declare no conflict of interest.

#### References

1. Fung, F.; Wang, H.S.; Menon, S. Food Safety in the 21st Century. *Biomed. J.* **2018**, *41*, 88–95. [[CrossRef](#)] [[PubMed](#)]
2. Chen, L.; Alali, W. Recent Discoveries in Human Serious Foodborne Pathogenic Bacteria: Resurgence, Pathogenesis, and Control Strategies. *Front. Microbiol.* **2018**, *9*, 2412. [[CrossRef](#)] [[PubMed](#)]
3. Xu, J.-G.; Cheng, B.-K.; Jing, H.-Q. *Escherichia Coli* O157: H7 and Shiga-like-toxin-producing *Escherichia Coli* in China. *World J. Gastroenterol.* **1999**, *5*, 191–194. [[CrossRef](#)] [[PubMed](#)]
4. Mylius, S.D.; Nauta, M.J.; Havelaar, A.H. Cross-contamination during food preparation: A mechanistic model applied to chicken-borne *Campylobacter*. *Risk Anal. Int. J.* **2007**, *27*, 803–813. [[CrossRef](#)] [[PubMed](#)]
5. Van Elsas, J.D.; Semenov, A.V.; Costa, R.; Trevors, J.T. Survival of *Escherichia coli* in the environment: Fundamental and public health aspects. *ISME J.* **2011**, *5*, 173–183. [[CrossRef](#)]
6. Waite, J.G.; Daeschel, M.A. Contribution of wine components to inactivation of food-borne pathogens. *J. Food Sci.* **2007**, *7*, M286–M291. [[CrossRef](#)]
7. Isohanni, P.; Alter, T.; Saris, P.; Lyhs, U. Wines as possible meat marinade ingredients possess antimicrobial potential against *Campylobacter*. *Poult. Sci.* **2010**, *89*, 2704–2710. [[CrossRef](#)] [[PubMed](#)]
8. Kargiotou, C.; Katsanidis, E.; Rhoades, J.; Kontominas, M.; Koutsoumanis, K. Efficacies of soy sauce and wine base marinades for controlling spoilage of raw beef. *Food Microbiol.* **2011**, *28*, 158–163. [[CrossRef](#)]
9. Tenore, G.C.; Basile, A.; Novellino, E. Antioxidant and antimicrobial properties of polyphenolic fractions from selected Moroccan red wines. *J. Food Sci.* **2011**, *76*, C1342–C1348. [[CrossRef](#)]
10. Papadopoulou, C.; Soulti, K.; Roussis, I.G. Potential antimicrobial activity of red and white wine phenolic extracts against strains of *Staphylococcus aureus*, *Escherichia coli* and *Candida albicans*. *Food Technol. Biotechnol.* **2005**, *43*, 41–46.
11. Friedman, M.; Henika, P.R.; Levin, C.E.; Mandrell, R.W. Antimicrobial wine formulations against the foodborne pathogens *Escherichia coli* O157:H7 and *Salmonella enterica*. *J. Food. Sci.* **2006**, *71*, M245–M251. [[CrossRef](#)]
12. Friedman, M. Antibacterial, antiviral, and antifungal properties of wines and winery byproducts in relation to their flavonoid content. *J. Agric. Food Chem.* **2014**, *62*, 6025–6042. [[CrossRef](#)] [[PubMed](#)]
13. Moretto, T.; Daeschel, M.A. Wine is bactericidal to food-borne pathogens. *J. Food Sci.* **2004**, *69*, M250–M257.

14. Radovanović, A.N.; Jovančičević, B.S.; Radovanović, B.C.; Mihajilov-Krstev, T. Antibacterial effectiveness of selected Vranac wines against six Gram-positive and six Gram-negative bacterial strains. *Trop. J. Pharm. Res.* **2014**, *13*, 819–824. [[CrossRef](#)]
15. Birk, T.; Knöchel, S. Fate of food-associated bacteria in pork as affected by marinade, temperature, and ultrasound. *J. Food Prot.* **2009**, *72*, 549–555. [[CrossRef](#)] [[PubMed](#)]
16. Nisiotou, A.; Chorianopoulos, N.G.; Gounadaki, A.; Panagou, E.Z.; Nychas, G.J. Effect of wine-based marinades on the behavior of *Salmonella Typhimurium* and background flora in beef fillets. *Int. J. Food Microbiol.* **2013**, *164*, 119–127. [[CrossRef](#)]
17. Rhoades, J.; Kargiotou, C.; Katsanidis, E.; Koutsoumanis, K.P. Use of marination for controlling *Salmonella enterica* and *Listeria monocytogenes* in raw beef. *Food Microbiol.* **2013**, *36*, 248–253. [[CrossRef](#)] [[PubMed](#)]
18. Díez, J.G.; Patarata, L. Behavior of *Salmonella* spp., *Listeria monocytogenes*, and *Staphylococcus aureus* in Chourico de Vinho, a dry fermented sausage made from wine-marinated meat. *J. Food Prot.* **2013**, *76*, 588–594. [[CrossRef](#)]
19. Friedman, M.; Levin, C.E.; Henika, P.R. Addition of phytochemical-rich plant extracts mitigate the antimicrobial activity of essential oil/wine mixtures against *Escherichia coli* O157:H7 but not against *Salmonella enterica*. *Food Control* **2017**, *73*, 562–565. [[CrossRef](#)]
20. Daglia, M. Polyphenols as antimicrobial agents. *Curr. Opin. Biotechnol.* **2012**, *23*, 174–181. [[CrossRef](#)]
21. Katalinić, V.; Možina, S.S.; Skroza, D.; Generalić, I.; Abramović, H.; Miloš, M.; Ljubenković, I.; Piskernik, S.; Pezo, I.; Terpinč, P.; et al. Polyphenolic profile, antioxidant properties and antimicrobial activity of grape skin extracts of 14 Vitisvinifera varieties grown in Dalmatia (Croatia). *Food Chem.* **2010**, *119*, 715–723. [[CrossRef](#)]
22. Milat, A.M.; Boban, M.; Teissedre, P.-L.; Šešelja-Perišin, A.; Jurić, D.; Skroza, D.; Generalić Mekinić, I.; Ljubenković, I.; Volarević, J.; Rasines-Perea, Z.; et al. Effects of oxidation and browning of macerated white wine on its antioxidant and direct vasodilatory activity. *J. Funct. Foods* **2019**, *15*, 138–147. [[CrossRef](#)]
23. Boban, N.; Tonkic, M.; Budimir, D.; Modun, D.; Sutlovic, D.; Punda-Polic, V.; Boban, M. Antimicrobial effects of wine: Separating the role of polyphenols, pH, ethanol, and other wine components. *J. Food Sci.* **2010**, *75*, M322–M326. [[CrossRef](#)]
24. Rodríguez-Vaquero, M.J.; Aredes-Fernández, P.A.; Manca de Nadra, M.C. Phenolic compounds from wine as natural preservatives of fish meat. *Food Technol. Biotechnol.* **2013**, *51*, 376–382.
25. Lee, J.H.; Cho, B.K. Effects of grape, wild grape and black raspberry wines on ground pork during refrigerated storage. *Acta Aliment.* **2014**, *43*, 553–563. [[CrossRef](#)]
26. Amerine, M.A.; Ough, C.S. *Methods for Analysis of Musts and Wines*, 1st ed.; John Wiley and Sons: New York, NY, USA, 1980; pp. 181–185.
27. Ribéreau-Gayon, P.; Stonestreet, E. Dosage des tanins du vin rouge et détermination de leur structure. *Chim. Anal.* **1966**, *48*, 188–196.
28. Ribéreau-Gayon, P.; Stonestreet, E. Le dosage des anthocyanes dans le vin rouge. *Bull. Soc. Chim. Fr.* **1965**, *9*, 2649–2652.
29. CLSI. *Methods for Dilution Antimicrobial Susceptibility Tests for Bacteria That Grow Aerobically*; Approved Standard—8th ed. CLSI document M07-A8; Clinical and Laboratory Standards Institute: Wayne, PA, USA, 2009; p. 226.
30. The European Committee on Antimicrobial Susceptibility Testing. Breakpoint Tables for Interpretation of MICs and Zone Diameters. Version 10.0, 2020. Available online: <http://www.eucast.org> (accessed on 7 May 2020).
31. CLSI. *Performance Standards for Antimicrobial Susceptibility Testing*; Twenty-Fourth Informational Supplement. CLSI document M100-S24; Clinical and Laboratory Standards Institute: Wayne, PA, USA, 2014.
32. Balouiri, M.; Sadiki, M.; Ibensoudaet, S.K. Methods for in vitro evaluating antimicrobial activity: A review. *J. Pharm. Anal.* **2016**, *6*, 71–79. [[CrossRef](#)]
33. Koohsari, H.; Ghaemi, E.A.; Shesh Poli, M.S.; Sadegh, A. Evaluation of antibacterial activity of *Lemon verbena* (*Lippia citriodora*) leaves. *Ann. Biol. Res.* **2013**, *4*, 52–55.
34. Skroza, D.; Šimat, V.; SmoleMožina, S.; Katalinić, V.; Boban, N.; Generalić Mekinić, I. Interactions of resveratrol with other phenolics and activity against food-borne pathogens. *Food Sci. Nutr.* **2019**, *7*, 2312–2318. [[CrossRef](#)]
35. ISO. *Microbiology of Food and Animal Feeding Stuffs—Horizontal Method for the Enumeration of Presumptive E. coli—Part 2: Colony Count Technique at 44 °C Using 5-Bromo-4-Chloro-3-Indolyl-Beta-D-Glucuronic Acid*; ISO 16649-2; International Organization for Standardization: Geneva, Switzerland, 2001; p. 8.

36. Šimat, V.; Bogdanović, T.; Krželj, M.; Soldo, A.; Maršić-Lučić, J. Differences in chemical, physical and sensory properties during shelf life assessment of wild and farmed gilthead sea bream (*Sparus aurata*, L.). *J. Appl. Ichtyol.* **2012**, *28*, 95–101. [CrossRef]
37. L 149/1 Official Journal of the European Union. Regulation (EU) No 1308/2013 of the European Parliament and of the Council as Regards Wine-Growing Areas where the Alcoholic Strength may be Increased, Authorised Oenological Practices and Restrictions Applicable to the Production and Conservation of Grapevine Products, the Minimum Percentage of Alcohol for By-Products and Their Disposal, and Publication of OIV Files. Available online: <https://eur-lex.europa.eu/legal-content/EN/TXT/PDF/?uri=OJ:L:2019:149:FULL&from=IT> (accessed on 15 March 2020).
38. Jamir, L.; Kumar, V.; Gaṭ, Y.; Kumar, A.; Kaur, S. Wine: A potential source of antimicrobial compounds. *J. Wine Res.* **2019**, *30*, 220–237. [CrossRef]
39. Just, J.R.; Daeschel, M.A. Antimicrobial effects of wine on *Escherichia coli* O157: H7 and *Salmonella typhimurium* in a model stomach system. *J. Food Sci.* **2003**, *68*, 285–290. [CrossRef]
40. Weisse, M.E.; Eberly, B.; Person, D.A. Wine as a digestive aid: Comparative antimicrobial effects of bismuth salicylate and red and white wine. *BMJ* **1995**, *311*, 1657–1660. [CrossRef] [PubMed]
41. Chan, M.M. Antimicrobial effect of resveratrol on dermatophytes and bacterial pathogens of the skin. *Biochem. Pharmacol.* **2002**, *63*, 99–104. [CrossRef]
42. Boban, N.; Tonkić, M.; Modun, D.; Budimir, D.; Mudnić, I.; Sutlović, D.; Punda-Polić, V.; Boban, M. Thermally treated wine retains antibacterial effects to food-borne pathogens. *Food Control* **2010**, *21*, 1161–1165. [CrossRef]
43. Papadopoulos, V.; Chouliara, I.; Badeka, A.; Savvaidis, I.N.; Kontominas, M.G. Effect of gutting on microbiological, chemical, and sensory properties of aquacultured sea bass (*Dicentrarchus labrax*) stored in ice. *Food Microbiol.* **2003**, *20*, 411–420. [CrossRef]
44. Maqsood, S.; Benjakul, S.; Shahidi, F. Emerging role of phenolic compounds as natural food additives in fish and fish products. *Crit. Rev. Food Sci. Nutr.* **2013**, *53*, 162–179. [CrossRef]
45. Cushnie, T.T.P.; Lamb, A.J. Recent advances in understanding the antibacterial properties of flavonoids. *Int. J. Antimicrob. Agents* **2011**, *38*, 99–107. [CrossRef]



© 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).